Supplementary information

The spontaneous location recognition task for assessing spatial pattern separation and memory across a delay in rats and mice

In the format provided by the authors and unedited

The Spontaneous Location Recognition Task for assessing spatial pattern separation and memory across a delay in rats and mice

Amy C. Reichelt ^{1,2,3,} Cecilia P. Kramar ⁴, Olivia R. Ghosh-Swaby ^{5,}
Paul A. S. Sheppard ^{1,2,} Brianne A. Kent ⁶, Pedro Bekinschtein ⁷,
Lisa M. Saksida ^{1,2,8,} Timothy J. Bussey ^{1,2,8,}

Corresponding authors

amy.reichelt@adelaide.edu.au, Adelaide Medical School, University of Adelaide, Adelaide, Australia

tbussey@uwo.ca, Translational Cognitive Neuroscience Laboratory, Western University, London, Ontario, Canada. <u>http://www.tcnlab.uwo.ca/</u>

Affiliations

1. Department of Physiology and Pharmacology, Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada

2. Robarts Research Institute, Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada

3. Adelaide Medical School, University of Adelaide, Adelaide, Australia

4. Douglas Hospital Research Center, Department of Psychiatry, McGill University, Montreal, Canada

5. Neuroscience Graduate Program, Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada

6. Department of Psychology, Simon Fraser University, Burnaby, British Columbia, Canada

7. Instituto de Neurociencia Cognitiva y Traslacional (CONICET-INECO-Universidad Favaloro), Buenos Aires, Argentina

8. Brain and Mind Institute, Western University, London, Ontario, Canada



Supplementary Figure 1. Examples of object location counterbalancing that can be used by a researcher who is running each condition twice for each of the SLR configurations. The order of the conditions (d-SLR, s-SLR, xs-SLR) should be counterbalanced across rats. Different sets of objects should be used in each of the tests.