

Title

Investigation of the pain factors of rotator cuff tears: a protocol for a scoping review

Authors

Tomohisa Yuda¹⁾, Takashi Saito²⁾, Hayato Shigetoh³⁾, Takashi Kitagawa⁴⁾

1. Department of Rehabilitation Medicine, Kitade Hospital, Wakayama, Japan

2. Department of Rehabilitation Medicine, Ono Orthopedic, Tochigi, Japan

3. Department of Physical Therapy, Faculty of Health Science, Kyoto Tachibana University, Kyoto, Japan

4. Department of Physical Therapy, School of Health Sciences, Shinshu University, Nagano, Japan

Abstract

- Objective : The purpose of this scoping review was to investigate the factors associated with rotator cuff tear pain and summarize them into the incidence, exacerbating, and remitting factors.
- Introduction : Rotator cuff tears are one of the most common shoulder disorders. A variety of associated factors have been cited in observational studies, and systematic reviews have examined factors associated with symptomatic rotator cuff tears. However, no reports differentiate pain factors into developmental, exacerbating, and remitting factors.
- Inclusion/exclusion criteria : The study will include patients whose physicians have diagnosed non-traumatic rotator cuff tears using Magnetic Resonance Imaging or ultrasound echo. Outcomes will be a pain, the severity of rotator cuff tear, physical function, and psychological factors. The study design will be observational, with no restrictions on region, race, gender, or language of the original paper.
- Methods : A systematic search of PubMed, Embase, Web of Science, Cumulative Index to Nursing and Allied Health Literature (CINAHL), and Physiotherapy Evidence Database (PEDro) databases using the keywords “rotator cuff tear”, “pain”, and “etiology” will be conducted during May 2022. In the first screening step, two independent reviewers will review all of the titles and abstracts to exclude irrelevant articles. In the second screening step, two independent reviewers will review all of the full texts to exclude irrelevant articles. Outcomes will focus on rotator cuff tear severity, pain, physical function, and psychosocial factors, categorizing factors associated with pain according to each study design and identifying incidence factors, exacerbating factors and remission factors.

Introduction

Rotator cuff tears are one of the most common shoulder joint disorders ¹⁾. Medical costs for arthroscopic rotator cuff repair in the U.S. are estimated to be \$1.2-1.6 billion annually, and these cost savings are expected to translate into lower medical costs ²⁾. Therefore, conservative treatment is an important stand-in, but it is not even clear why the pain associated with rotator cuff tears occurs.

Cross-sectional studies have reported that the presence of pain and weakness in shoulder abduction and external rotation, impingement sign, and dominant arm are factors associated with symptomatic rotator cuff tears, but not the severity of rotator cuff tear ³⁻⁴⁾. Prospective studies examining the developmental factors in patients with asymptomatic rotator cuff tears have identified increased tear size in full-thickness tears ⁵⁻⁷⁾, a transition from partial to full tears ⁵⁻⁷⁾, development of lesions in the long head of the biceps brachii muscle ⁷⁾, and fatty degenerative ⁷⁾, as factors in the development of pain. A prospective study investigating pain transitions in symptomatic rotator cuff tears found that enlargement of the rotator cuff tear was not associated with pain intensity ⁸⁻⁹⁾.

Thus, it can be inferred that factors associated with rotator cuff tear pain differ depending on the study design and subject. Therefore, it can be inferred that factors associated with rotator cuff tear pain vary by study design and subject. Therefore, we aimed to summarize factors associated with rotator cuff tear pain by distinguishing between relevant factors from cross-sectional studies and factors associated with pain development, exacerbation, and remission from prospective studies.

Review Question

The purpose of this scoping review is to investigate the factors associated with pain in degenerative rotator cuff tears and to summarize development, exacerbation, and remission factors.

Keywords

Pain, Risk factors, Rotator cuff tear, Shoulder, Symptomatic

Eligibility Criteria

Patients will be included in the physician diagnoses a rotator cuff tear using Magnetic Resonance Imaging or ultrasound echocardiography. Exclusion criteria for the study include those with traumatic episodes, unmeasurable partial or extensive tears, neurological deficits, surgical intervention, and shoulder joint disease other than rotator cuff tears.

Concept

【Patient】

- Degenerative rotator cuff tear (Non-traumatic)

【Exposure】

- Conservative therapy, rehabilitation, physical therapy, drug therapy, injection therapy

【Outcome】

- Does not select specific evaluation items but use terms for relevant factors

【Study design】

- Observational studies: cross-sectional studies, prospective cohort studies, retrospective cohort studies, case-control studies

Context

The search results are limited to original papers published in peer-reviewed journals, with no language restrictions. The publication dates shall be through April 2022.

Types of Sources

Observational studies will be included. Interventional trials and systematic reviews will not be included. Intervention studies including randomized controlled trials (RCT) and non-RCT, review papers such as randomized reviews, systematic reviews, and meta-analysis, as well as case reports and case series studies that mainly report intervention effects will be excluded from this study.

Methods

This protocol was developed based on PRISMA 2020 statement. The scoping review will also be conducted based on the scoping review methodology by the JBI.

Search Strategy

The search strategy will be a systematic electronic search using the following databases with the aim of finding published studies: PubMed, Embase, Web of Science, CINAHL, and PEDro. A complete search strategy for the five databases will be developed using the words in the titles and abstracts of the relevant articles (the appendix contains more details).

Study/ Source of Evidence Selection

After the search, all identified citations will be collated and uploaded to Rayyan to remove duplicates. In a pilot test, the title and abstract will be reviewed by two independent reviewers

and evaluated against the review's inclusion criteria. All potentially relevant papers will be searched, and details of their citations will be incorporated into Rayyan. The full text of the selected citations will be evaluated in detail by two independent reviewers based on the inclusion criteria. The scoping review will note and report the reasons for excluding full-text references that do not meet the inclusion criteria. Disagreements that arise among reviewers at each stage of the selection process will be discussed or resolved with additional reviewers. Search results and the study uptake process will be reported at the final scoping review and presented using the PRISMA 2020 statement flow diagram for scoping reviews.

Data Extraction

Data extraction will be performed using a data extraction tool. Data will specifically show findings related to participants, concepts, context, study design, and review questions. Extracted forms will include information such as "author," "title," "year of publication," "study design," "language" "study population characteristics," "independent variable outcomes," and "dependent variable outcomes."

The draft data extraction will be modified and revised as needed during the process of extracting data from each of the included evidence sources. The revisions will be described in the scoping review.

Disagreements arising among reviewers will be resolved through discussion or with additional reviewers. Incomplete reporting of critical data will be treated as missing data if no response is received within one month of the e-mail communication.

Data Analysis and Presentation

The data will be presented in graphical or illustrative or tabular format, depending on the study design and risk factors.

Acknowledgments

None.

Funding

None.

Conflicts of Interest

None.

References

1. Jeong JJ, Park SE, Ji JH, Lee HH, Jung SH, Choi BS. Trans-tendon suture bridge rotator cuff repair with tenotomized pathologic biceps tendon augmentation in high-grade PASTA lesions. *Arch Orthop Trauma Surg.* 2020;140(1):67-76.
2. Chalmers PN, Granger E, Nelson R, Yoo M, Tashjian RZ. Factors affecting cost, outcomes, and tendon healing after arthroscopic rotator cuff repair. *Arthroscopy: The Journal of Arthroscopic & Related Surgery.* 2018;34(5):1393-1400.
3. Yamamoto A, Takagishi K, Kobayashi T, Shitara H, Osawa T. Factors involved in the presence of symptoms associated with rotator cuff tears: a comparison of asymptomatic and symptomatic rotator cuff tears in the general population. *Journal of Shoulder and Elbow Surgery.* 2011;20(7):1133-1137.
4. Dunn WR, Kuhn JE, Sanders R, et al. Symptoms of pain do not correlate with rotator cuff tear severity: a cross-sectional study of 393 patients with a symptomatic atraumatic full-thickness rotator cuff tear. *Journal of Bone and Joint Surgery.* 2014;96(10):793-800.
5. Mall NA, Kim HM, Keener JD, et al. Symptomatic progression of asymptomatic rotator cuff tears. *J Bone Joint Surg Am.* 2010;92(16):2623-2633.
6. Keener JD, Galatz LM, Teefey SA, et al. A prospective evaluation of survivorship of asymptomatic degenerative rotator cuff tears. *J Bone Joint Surg Am.* 2015;97(2):89-98.
7. Moosmayer S, Tariq R, Stiris M, Smith HJ. The natural history of asymptomatic rotator cuff tears: a three-year follow-up of fifty cases. *The Journal of Bone and Joint Surgery-American Volume.* 2013;95(14):1249-1255.
8. Yamamoto N, Mineta M, Kawakami J, Sano H, Itoi E. Risk factors for tear progression in symptomatic rotator cuff tears: a prospective study of 174 shoulders. *Am J Sports Med.* 2017;45(11):2524-2531.
9. Kim YS, Kim SE, Bae SH, Lee HJ, Jee WH, Park CK. Tear progression of symptomatic full-thickness and partial-thickness rotator cuff tears as measured by repeated MRI. *Knee Surg Sports Traumatol Arthrosc.* 2017;25(7):2073-2080.

Appendix: Search strategy

☐ PubMed search strategy

("rotator cuff injuries"[MeSH Terms] OR "rotator cuff tear*" [Title/Abstract] OR "rotator cuff injur*" [Title/Abstract] OR "rotator cuff tendin*" [Title/Abstract] OR "rotator cuff related pain" [Title/Abstract]) AND ("association"[MeSH Terms] OR "associat*" [Title/Abstract] OR "risk"[MeSH Terms] OR "risk" [Title/Abstract] OR "causality"[MeSH Terms] OR "etiology" [Title/Abstract] OR "pathology"[MeSH Terms] OR "pathology" [Title/Abstract] OR "cause" [Title/Abstract] OR "predict*" [Title/Abstract]) AND ("pain"[MeSH Terms] OR "pain" [Title/Abstract] OR "symptomatic" [Title/Abstract] OR "asymptomatic" [Title/Abstract])

☐ Embase

(MESH.EXACT.EXPLODE("rotator cuff injuries") OR TI,AB("rotator cuff tear*") OR TI,AB("rotator cuff injur*") OR TI,AB("rotator cuff tendin*") OR TI,AB("rotator cuff related pain")) AND (MESH.EXACT.EXPLODE(association) OR TI,AB(associat*) OR MESH.EXACT.EXPLODE(risk) OR TI,AB(risk) OR MESH.EXACT.EXPLODE(causality) OR TI,AB(etiology) OR MESH.EXACT.EXPLODE(pathology) OR TI,AB(pathology) OR TI,AB(cause) OR TI,AB(predict*)) AND (MESH.EXACT.EXPLODE(pain) OR TI,AB(pain) OR TI,AB(symptomatic) OR TI,AB(asymptomatic))

☐ Web of science search strategy

- ① rotator cuff injuries OR rotator cuff tear* OR rotator cuff injur* OR rotator cuff tendin* OR rotator cuff related pain)
- ② associat* OR risk OR causality OR etiology OR pathology OR cause OR predict*
- ③ pain OR pain OR symptomatic OR asymptomatic
- ④ ① and ② and ③

☐ CINAHL search strategy

((MH "rotator cuff injuries"+) OR (TI "rotator cuff tear*" OR AB "rotator cuff tear*") OR (TI "rotator cuff injur*" OR AB "rotator cuff injur*") OR (TI "rotator cuff tendin*" OR AB "rotator cuff tendin*") OR (TI "rotator cuff related pain" OR AB "rotator cuff related pain")) AND ((MH association+) OR (TI associat* OR AB associat*) OR (MH risk+) OR (TI risk OR AB risk) OR (MH causality+) OR (TI etiology OR AB etiology) OR (MH pathology+) OR (TI pathology OR AB pathology) OR (TI cause OR AB cause) OR (TI predict* OR AB predict*)) AND ((MH pain+) OR (TI pain OR AB pain) OR (TI symptomatic OR AB symptomatic) OR (TI asymptomatic OR AB asymptomatic))

☐ Physiotherapy Evidence Database (PEDro) Search strategy

Abstract & Title: rotator and cuff*